



## Burglind Juhl-Jöricke

### Analytic knots, satellites and the 4-ball genus

**ZEIT:**

6.7.2012, 10:00 Uhr - 11:00 Uhr

**ORT:**

HU  
Institut für Mathematik, Raum 1.114  
Rudower Chaussee 25  
12489 Berlin

Abstract. A satellite of a knot  $K$  is a link contained in a tubular neighbourhood of  $K$  (but not in a 3-ball contained in this neighbourhood), and not isotopic to  $K$ . In a classical paper Schubert gives a lower bound for the genus of satellites.

Consider a knot or link in the unit sphere in  $\mathbb{C}^2$ . Call it analytic (respectively, smoothly analytic) if it bounds a complex curve (respectively, a smooth complex curve) in the complex ball.

Let  $K$  be a smoothly analytic knot. For analytic satellite links contained in a sufficiently small tubular neighbourhood of  $K$  there is a (sharp) lower bound of the 4-ball genus (but not of the genus).

Moreover, these links can be completely described. The problem is related to branched coverings and braided surfaces.

**Kontakt:**

Humboldt-Universität zu Berlin . Institut für Mathematik  
SFB 647 . Unter den Linden 6 . 10099 Berlin  
Tel. +49 30 2093 1804 . Fax. +49 30 2093 2727  
sfb647@math.hu-berlin.de

[www.raumzeitmaterie.de](http://www.raumzeitmaterie.de)